
ACCEPTED FOR REGISTRATION

JULY 07 2016

New York State Department of Environmental Conservation Division of Materials Management Pesticide Product Registration

Doc ID: 545807



For Up To 4 Months

0 7 5



TOP! READ ENTIRE LABEL BEFORE USE. DIRECTIONS FOR USE It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

How It Works
The Fruit Fly BarPro Fly Strip utilizes controlled release technology to slowly diffuse a deep penetrating vapor in enclosed spaces for up to 4 months. The clean, odorless vapor is evenly distributed throughout the enclosed treatment area, killing visible and hidden insects on contact and preventing new insect infestations (while you're away).

What It Does
Kills flying and crawling insects including: flies, gnats, mosquitoes, moths, silverfish, cockroaches, spiders, beetles, earwigs, spider mites and other listed insects.

Where To Use Garbage rooms, lavatories, floor drains (to sewer), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, utility equipment, garages, mop closets, and product storage, after canning or bottling.

RESTRICTIONS

- Do not use in the food/feed areas of food/feed processing or food/feed manufacturing or food/feed service establishments. Non-food/feed areas of these establishments where the product can be used are areas such as garbage rooms, lavatories, floor drains (to sewer), entries and vestibules, offices, locker rooms, machine rooms, boiler rooms, utility equipment, garages, mop closets, and product storage (after canning or hottling).

Storage (after carning or botting).
If product is used within homes, use only in closets, wardrobes and cupboards.
Do not use in areas where people will be for an extended period of time.
Do not allow children to handle new or used strips.

This product controls insects as listed. This product may be used in non-food/feed areas of food/feed processing or food/feed manufacturing or food/feed service establishments. Remove pre-assembled pressing or food/feed manufacturing or food/feed service establishments. Remove pre-assembled pressing from the sealed bag when ready to use. Do not allow children to handle new or used strips. Place in the desired location away from windows and out of the reach of children and pets. Drafts, weather and other conditions may affect the performance, but treatment usually lasts for four months. Record the date of installation and replace with a new strip at the end of four months or when effectiveness diminishes. Always use the appropriate size and number of strips for the space being treated. One 16g strip will treat 200 cu. ft. If more than one strip is needed for the space being treated, space strips evenly throughout the treatment area. In non-food/feed areas of food/feed processing or food/feed manufacturing or food/feed areas of food/feed areas

STORAGE AND DISPOSAL
Storage: Do not contaminate water, food or feed by storage or disposal. Do not open or remove from package before ready to install. Store in a cool, dry place.

Polich Do not open or remove from package before ready to use, **Disposal: If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain. **Outer Container**Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available or dispose of the empty container in trash.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be fatal if swallowed. Do not get in mouth. Harmful if inhaled. Causes moderate eye irritation. Avoid breathing vapors. Avoid contact with eyes or clothing. After prolonged storage, a small amount of liquid may form on strip. Do not get liquid in eyes. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove and wash contaminated clothing before reuse. Do not use in hospitals or clinic rooms, such as patient rooms, wards, nurseries, operating or emergency areas. If used within homes, use only in closets, wardrobes and cupboards, but do not use in closets and wardrobes of rooms where infants, children and the sick or aged are or will be present for any extended period of confinement. Do not use in any area of the home where people will be present for extended periods of time. Do not use in kitchens, except cupboards. Do not use in the food/feed area of restaurants or any area where food is prepared or served. Not to be taken internally by humans or animals.

FIRST AID

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor. Do not give anything by mouth to an unconscious person.


IF INHALED: Move person to fresh air, If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing, Rinse skin immediately with plenty of water for 15-20 minutes.
Call a poison control center or doctor for treatment advice.

Call a poison control center or doctor for treatment advice.

NOTE TO PHSYCIAN: Probable mucosal damage may contraindicate the use of gastric lavage. POISONING SYMPTOMS: CONTAINS AN ORGANOPHOSPHATE THAT INHIBITS CHOLINESTERASE. Symptoms include weakness, headache, tightness in chest, blurred vision, non-reactive pin point pupils, salivation, sweating, nausea, vomiting, diarrhea, and abdominal cramps. TREATMENT: Atropine is the specific therapeutic antagonist of choice against parasympathetic nervous stimulation. If there are signs of parasympathetic stimulation, atropine sulfate should be injected at 10 minute intervals in doses of 1 to 2 milligrams until complete atropinization has occurred. Morphine is contraindicated. 2-PAM is also antidotal and may be administered in conjunction with atropine. Clear chest by postural drainage.



Have the product container or label with you when calling a poison control center or doctor, or going for treatment. FOR THE FOLLOWING EMERGENCIES, PHONE 24 HOURS A DAY:

www.FruitFlyBarPro.com EPA Reg. No. 5481-348-87714 EPA Est. No. 42836-TX-1