DOC ID: 543314

## ACCEPTED FOR REGISTRATION

September 18, 2015

New York State Department of Environmental Conservation Division of Materials Management Pesticide Product Registration TEAR TOWARD CENTER FROM BOTH NOTCHES PRESS ZIPPER TO RESEAL 12455-79

Inner Label Prints at 100% on 8.5x11 Page 1 of 2

# KILLS RATS, MICE, AND MEADOW VOLES\*

**ALL-WEATHER BLOX™** 

# **Kills Warfarin Resistant Norway Rats**





12455-79

Inner Label Prints at 100% on 8.5x11 Page 2 of 2



# **KILLS RATS, MICE, AND MEADOW VOLES\***

### FIRST AID

#### HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE IF SWALLOWED:

- Call a poison control center, doctor, or 1-877-854-2494, or 1-800-858-7378\*\* immediately for treatment advice.
- Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.
- IF ON SKIN OR CLOTHING:
- Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes
- · Call a poison control center or doctor for treatment advice.
- IF IN EYES: · Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye
- · Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice
- \*\* Also call this number for information on health concerns and pesticide incidents

### NOTE TO PHYSICIAN

If swallowed or absorbed through the skin, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K1 intramuscularly or orally Repeat as necessary based on monitoring of prothrombin times.

### TREATMENT FOR PET POISONING If animal eats bait, call veterinarian at once

### NOTE TO VETERINARIAN

Anticoagulant Bromadiolone: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K1. If needed, check prothrombin times every 3 days until values return to normal (up to 30 days). In severe cases, blood transfusions may be needed.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for sites, pests, and application methods described on this label.

IMPORTANT: Do not expose children, pets, or nontarget animals to rodenticides. To help to prevent exposure:

- 1. Store unused product out of reach of children and pets.
- 2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
- 3. Dispose of product container and unused, spoiled, or unconsumed bait as specified on this label

#### Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used wherever children, pets, non-target mammals, or birds may have access to the bait placement location.

USE RESTRICTIONS: This product may only be used to control the following rodent pests in and around man-made structures: House mouse (Mus musculus), Norway rat (Rattus norvegicus), Roof rat (Rattus rattus), Cotton mouse (Peromyscus gossypinus), Cotton rat\* (Sigmodon hispidus), Deer mouse (Peromysus maniculatus), Eastern harvest mouse\* (Reithrodontomys humuli), Golden mouse\* (Ochrotomys nuttalli), Polynesian rat\* (Rattus exulans), Meadow vole (Microtus pennsylvanicus), White-footed mouse (Peromyscus leucopus), White-throated woodrat\* (Neotoma albigula), Southern plains woodrat\* (Neotoma micropus), and Mexican

### Kills Warfarin Resistant Norway Rats

Norway rats, roof rats, and house mice may consume a lethal dose in one night's feeding with first dead rodents appearing four or five days after feeding begins.

ACTIVE INGREDIENT:

Bromadiolone (CAS #28772-56-7): . . . . . . . . 0.005% tContains Denatonium Benzoate . 99.995% TOTAL

# **KEEP OUT OF REACH OF CHILDREN** CAUTION

See side panels for First Aid and additional precautionary statements.

### ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish, birds and other wildlife. Dogs and predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff also may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or rinsate

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Pesticide Storage: Store only in original container in a cool, dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

Pesticide Disposal: Wastes resulting from the use of this product may be placed in trash or delivered to an approved waste disposal facility.

Container Handling: Nonrefillable container. Do not reuse or refill this container. [Plastic:] Offer for recycling or reconditioning; or puncture and dispose of in a sanitary landfill; or by incineration. In most states, burning is not allowed.

WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions

## NET WEIGHT: 1 Ib (0.45 kg)

### INDIVIDUAL SALE IS PROHIBITED BY LAW

EPA REG. NO. 12455-79

EPA EST. NO. 12455-WI-1



Madison, WI 53704 U.S.A www.belllabs.com

#### DIRECTIONS FOR USE (Continued from other panel)

woodrat\* (Neotoma mexicana). This product must be used in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops.

Burrow baiting with Contrac All-Weather Blox is prohibited.

Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations. Do not broadcast bait. Do not use this product in sewers

Do not sell this product in individual containers holding less than 16 pounds of bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats, mice, or meadow voles\* will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in corners and concealed places, between floors and walls, or in locations where rats. mice, or meadow voles\*, or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

#### APPLICATION DIRECTIONS:

RATS: Place 3 to 16 bait blocks (at intervals of 15 to 30 feet) per placement in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

MICE AND MEADOW VOLES\*: Place 1 block per placement. Space placements at 8- to 12-foot intervals in infested areas. Two blocks may be needed at points of very high activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse or meadow vole\* activity cease

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Wearing gloves, collect and dispose of all dead, exposed animals and leftover bait. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through the skin. Keep away from children, domestic animals and pets. Do not get in eyes, on skin or on clothing.

All handlers (including applicators) must wear: shoes plus socks, and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves

### User Safety Requirements

Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

\*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse. Golden mouse. Polynesian rat. Meadow vole. White-throated woodrat. Southern plains woodrat, and Mexican woodrat

# 3699 Kinsman Blvd.

## DOC ID: 543314

### FIRST AID HAVE LABEL WITH YOU WHEN OBTAINING TREATMENT ADVICE

### IF SWALLOWED:

· Call a poison control center, doctor, or 1-877-854-2494, or 1-800-858-7378\*\* immediately for treatment advice

· Have person sip a glass of water if able to swallow.

Do not induce vomiting unless told to do so by the poison control center or doctor.

### IF ON SKIN OR CLOTHING:

 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15–20 minutes. Call a poison control center or doctor for treatment advice.

### IF IN FYES

- · Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- · Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center, doctor, or 1-877-854-2494 immediately for treatment advice.

\*\* Also call this number for information on health concerns and pesticide incidents.

### NOTE TO PHYSICIAN

If swallowed or absorbed through the skin, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K1 intramuscularly or orally. Repeat as necessary based on monitoring of prothrombin times.

# TREATMENT FOR PET POISONING

If animal eats bait call veterinarian at once NOTE TO VETERINARIAN

Anticoagulant Bromadiolone: For animals ingesting bait and/or showing poisoning signs (bleeding or elevated prothrombin times), give Vitamin K1. If needed, check prothrombin times every 3 days until values ACCEPTED return to normal (up to 30 days). In severe cases, blood transfusions may be needed. FOR REGISTRATION

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. Use only for sites, New York State Department

of Environmental Conservation See side panels for First Aid and additional precautionary statements. IMPORTANT: Do not expose children, pets, or nontarget animals to rodenticides. To help to prevent exposure: Division of Materials Management

September 18, 2015

2. Apply bait in locations out of reach of children, pets, domestic animals and nontargetpieleffeefdee Product Registration tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six vears of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from bait stations when they are lifted, units must be secured or otherwise immobilized. Stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.

3. Dispose of product container and unused, spoiled, or unconsumed bait as specified on this label.

### Bait stations are mandatory for outdoor, above-ground use. Tamper-resistant bait stations must be used wherever children, pets, non-target mammals, or birds may have access to the bait placement location.

USE RESTRICTIONS: This product may only be used to control the following rodent pests in and around man-made structures: House mouse (Mus musculus), Norway rat (Rattus norvegicus), Roof rat (Rattus rattus), Cotton mouse (Peromyscus gossypinus), Cotton rat\* (Sigmodon hispidus), Deer mouse (Peromysus maniculatus), Eastern harvest mouse\* (Reithrodontomys humuli), Golden mouse\* (Ochrotomys nuttalli), Polynesian rat\* (Rattus exulans), Meadow vole\* (Microtus pennsylvanicus), White-footed mouse (Peromyscus leucopus), White-throated woodrat\* (Neotoma albigula), Southern plains woodrat\* (Neotoma micropus), and Mexican woodrat\* (Neotoma mexicana). This product must be used in and within 100 feet of man-made structures constructed in a manner so as to be vulnerable to commensal rodent invasions and/or to harboring or attracting rodent infestations. Examples of such structures include homes and other permanent or temporary residences, food processing facilities, industrial and commercial buildings, trash receptacles, agricultural and public buildings, transport vehicles (ships, trains, aircraft), docks and port or terminal buildings and related structures around and associated with these sites. Fence and perimeter baiting, beyond 100 feet from a structure as defined above, is prohibited. This product must not be applied directly to food or feed crops.

Burrow baiting with Contrac All-Weather Blox is prohibited.

Do not place near or inside ventilation duct openings. Do not contaminate water, food, feedstuffs, food or feed handling equipment, or milk or meat handling equipment or surfaces that come into direct contact with food. When used in USDA inspected facilities, this product must be applied in tamper-resistant bait stations. Do not broadcast bait. Do not use this product in sewers.

# CONTRAC ALL-WEATHER BLOX™

### **KILLS RATS, MICE, AND MEADOW VOLES\***

### **Kills Warfarin Resistant Norway Rats**

Norway rats, roof rats, and house mice may consume a lethal dose in one night's feeding with first dead rodents appearing four or five days after feeding begins.

ACTIVE INGREDIENT:		
Bromadiolone (CAS #28772-56-7):		
OTHER INGREDIENTS <sup>†</sup> :		. 99.995%
E Contains Denatonium Benzoate	TOTAL	100.000%

KEEP OUT OF REACH OF CHILDREN

CAUTION

STORAGE AND DISPOSAL

Pesticide Storage: Store only in original container in a cool, dry place inaccessible to children and pets. Keep

Pesticide Disposal: Wastes resulting from the use of this product may be placed in trash or delivered to an

Container Handling: Nonrefillable container. Do not reuse or refill this container. [Plastic:] Offer for recycling or

reconditioning; or puncture and dispose of in a sanitary landfill; or by incineration. In most states, burning is not

allowed. [Paper:] Dispose of empty container by placing in trash, at an approved waste disposal facility or by

DIRECTIONS FOR USE (Continued from other panel)

Do not sell this product in individual containers holding less than 16 pounds of bait.

SELECTION OF TREATMENT AREAS: Determine areas where rats, mice, or meadow voles\* will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in corners and concealed places, between floors and walls, or in locations where rats, mice, or meadow voles\*, or their signs have been seen. Protect bait from rain and snow. Remove as much alternative food as possible.

### APPLICATION DIRECTIONS:

RATS: Place 3 to 16 bait blocks (at intervals of 15 to 30 feet) per placement in infested areas. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of rat activity cease.

MICE AND MEADOW VOLES\*: Place 1 block per placement. Space placements at 8- to 12-foot intervals in infested areas. Two blocks may be needed at points of very high activity. Maintain an uninterrupted supply of fresh bait for at least 15 days or until signs of mouse or meadow vole\* activity cease.

FOLLOW-UP: Replace contaminated or spoiled bait immediately. Wearing gloves, collect and dispose of all dead, exposed animals and leftover bait. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish as needed.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed or absorbed through the skin. Keep away from children, domestic animals and pets. Do not get in eves, on skin or on clothing.

All handlers (including applicators) must wear: shoes plus socks, and waterproof gloves. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

### **User Safety Requirements**

Follow manufacturer's instruction for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash hands thoroughly after applying bait and before eating, drinking, chewing gum, using tobacco or using the toilet and change into clean clothing.

### ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish, birds and other wildlife. Dogs and predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not apply this product directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff also may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash water or rinsate.

\*Not permitted for use against the following species in California: Cotton rat, Eastern harvest mouse, Golden mouse, Polynesian rat, Meadow vole, White-throated woodrat, Southern plains woodrat, and Mexican woodrat

090415/09-15

WARRANTY: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

# NET WEIGHT: 16 lbs (7.2 kg)

EPA REG. NO. 12455-79

EPA EST. NO. 12455-WI-1



Do not contaminate water, food or feed by storage or disposal.

containers closed and away from other chemicals.

incineration. In most states, burning is not allowed.

approved waste disposal facility.

3699 Kinsman Blvd Madison, WI 53704 U.S.A. www.belllabs.com MADE IN USA



12455-79 Outer Label Prints at 100% on 8.5x11